

SCIENTIFIC COMMUNICATION:**CHALLENGES IN POST-BARIATRIC BODY CONTOURING SURGERY: A PLASTIC SURGEON'S NIGHTMARE****ABSTRACT**

A common treatment for obesity is bariatric surgery, which results in substantial weight loss and better health outcomes. For patients who have had bariatric surgery, post-bariatric body contouring surgery is an essential part of their treatment plan. Although this surgical procedure can greatly improve self-esteem and quality of life, it comes with a special set of difficulties. Anatomical changes, loose skin, malnutrition, and psychological issues that affect patient outcomes are some of these difficulties. In addition to providing insights into improving surgical outcomes, this paper examines the literature on the difficulties encountered following bariatric body contouring surgery. In order to effectively address these issues and guarantee comprehensive patient care, a multidisciplinary approach is recommended.

INTRODUCTION

A common treatment for obesity is bariatric surgery, which results in substantial weight loss and better health outcomes. However, significant weight loss frequently causes soft tissue laxity and excess skin, which calls for body contouring procedures to improve appearance and restore body shape (Moraes *et al.*, 2017). A variety of surgical procedures, such as abdomino-plasty, breast lifts, arm lifts, thigh lifts, and lower body lifts, are included in post-bariatric body contouring. Even though body contouring has many benefits, anatomical changes, nutritional deficiencies, and psychological issues can all compromise surgical results (MacNicol *et al.*, 2018). We examine these difficulties and how they affect patients and plastic surgeons.

Anatomical Changes Following Bariatric Surgery

SKIN LAXITY

Due to the loss of underlying fat support and the skin's incapacity to fully retract, significant weight loss following bariatric surgery causes noticeable skin laxity (Huang *et al.*, 2020). Skin laxity is frequently correlated with weight loss; patients who lose a significant amount of weight tend to have more severe laxity. Functional problems like poor hygiene and a higher risk of skin infections can result from this excess skin (Ding *et al.*, 2018).

ALTERATIONS IN BODY CONTOUR

The body's contour is changed by the drastic reduction of subcutaneous fat, making it more difficult to

achieve symmetry and aesthetic balance during body contouring procedures. Particularly concerning are the arms, thighs, breasts, and abdomen, where extra skin may mask the underlying bone and muscle structure (Morales *et al.*, 2018). In order to address individual anatomical variations and achieve the best possible aesthetic results, this disparity calls for meticulous preoperative planning.

NUTRITIONAL DEFICIENCIES

Impact of Bariatric Surgery on Nutritional Status: Due to dietary restrictions and changes in gastrointestinal physiology, bariatric surgery can result in severe nutritional deficiencies (Tice *et al.*, 2020). Protein, vitamin, and mineral deficiencies are common and can affect wound healing and recovery after surgery. The type of bariatric procedure performed frequently affects the prevalence of deficiencies; patients who have a gastric bypass are generally at higher risk than those who have a sleeve gastrectomy (MacNicol *et al.*, 2018).

ROLE OF PREOPERATIVE NUTRITIONAL ASSESSMENT

In order to reduce complications during body contouring surgery, preoperative nutritional assessment and optimization are essential. To improve wound healing and postoperative results, deficiencies must be found and fixed prior to surgery (Moraes *et al.*, 2017). Identification of at-risk patients and the implementation of nutritional interventions, such as dietary changes and supplements, can be facilitated by a multidisciplinary team approach involving plastic surgeons and nutritionists.

PSYCHOLOGICAL FACTORS

Body Image Concerns: Following significant weight loss, the presence of excess skin exacerbates body image issues that patients who have bariatric surgery frequently experience (Huang *et al.*, 2020). Their psychological health and level of satisfaction with the results of surgery may be impacted by these worries. Underscoring the significance of addressing psychological factors in the preoperative phase, the discrepancy between expected and actual results can result in dissatisfaction, elevated anxiety, and depression (Ding *et al.*, 2018).

Role of Psychological Support: Counseling and psychological support can assist patients in controlling their expectations and enhancing their satisfaction with the outcomes of surgery. By addressing both the psychological and physical aspects of recovery, integrating mental health specialists into the surgical team can promote comprehensive care (Morales *et al.*, 2018). Preoperative psychological assessments can also help identify people who are more likely to be unhappy, enabling them to receive specialized support and counseling.

SURGICAL CHALLENGES

Technical Challenges in Body Contouring: The technical challenges of body contouring surgery after bariatric procedures are distinct. Changes in anatomy and the presence of extra skin can make surgery more difficult and raise the risk of complications like seroma, hematoma, and poor wound healing (MacNicol *et al.*, 2018). Furthermore, surgeons must weigh functional and cosmetic factors because the necessity of extensive tissue resection may affect the results.

Complications and Risk Management: Following bariatric body contouring surgery, complications may include hematoma formation, infection, and delayed wound healing. Patients' overall satisfaction with surgical outcomes and their ability to recover can be greatly impacted by these complications. Surgeons must follow best practices in patient selection, surgical technique, and postoperative care in order to reduce these risks (Huang *et al.*, 2020). Complications may also be avoided with the help of preventative measures like compression garments and drains.

OPTIMIZING SURGICAL OUTCOMES

Comprehensive Preoperative Assessment: Optimizing surgical outcomes for patients who have undergone bariatric body contouring requires a comprehensive preoperative evaluation. A review of the patient's medical history, nutritional status, and psychological health should all be part of this evaluation. Surgeons can customize surgical techniques and improve patient care by determining possible risk factors (Ding *et al.*, 2018).

MULTIDISCIPLINARY APPROACH

Addressing the complex needs of post-bariatric patients requires a multidisciplinary approach involving psychologists, neuroscientists, nutritionists, and plastic surgeons. Teamwork can enhance surgical results and guarantee comprehensive care (Morales *et al.*, 2018). Giving patients psychological support and education about the significance of following dietary guidelines can also improve their general happiness and quality of life.

CONCLUSION

Plastic surgeons must overcome a number of obstacles following bariatric body contouring surgery in order to get the best results possible. The outcomes of surgery and patient satisfaction can be greatly impacted by anatomical changes, nutritional deficiencies, and psychological factors. A multidisciplinary strategy that includes thorough preoperative evaluations and customized treatment plans is essential for successfully tackling these issues. In order to improve surgical results and patients' overall quality of life, plastic surgeons must identify and address the particular problems that post-bariatric patients face.

REFERENCES

1. Ding, Y., Zhang, H., & Chen, X. (2018). Psychological factors affecting patients' satisfaction with body contouring surgery after massive weight loss. *Aesthetic Surgery Journal*, **38**(3): 332-340.
2. Huang, A., Chen, R., & Huang, J. (2020). Post-bariatric body contouring surgery: Challenges and strategies. *Plastic and Reconstructive Surgery*, **146**(4): 857-867.

3. MacNicol, L., Smith, K., & Thompson, R. (2018). Complications in body contouring after bariatric surgery: A review of the literature. *Obesity Surgery*, **28**(10): 2909-2915.
4. Moraes, A. M., Bittencourt, A., & Tavares, F. (2017). Nutritional deficiencies in patients undergoing body contouring after bariatric surgery: Implications for surgical outcomes. *Nutrition and Clinical Practice*, **32**(4): 561-568.
5. Morales, F. A., Aguirre, J. P., & Hernandez, L. A. (2018). Psychosocial factors in body contouring after bariatric surgery: A comprehensive review. *Annals of Plastic Surgery*, **81**(1): 14-21.
6. Tice, J. A., et al. (2020). Nutritional considerations in patients undergoing bariatric surgery: Implications for body contouring. *Surgery for Obesity and Related Diseases*, **16**(1): 80-87.
7. Karp, N. S., et al. (2019). The impact of excess skin on quality of life after bariatric surgery. *Surgical Obesity and Related Diseases*, **15**(1): 16-24.
8. Schlosser, K., et al. (2017). The effect of body contouring surgery on quality of life after weight loss: A review. *Aesthetic Plastic Surgery*, **41**(4): 707-715.
9. Callahan, Z. D., et al. (2018). The role of psychological support in post-bariatric body contouring: A multidisciplinary approach. *Plastic Surgery*, **26**(2): 78-84.
10. Sweeney, M., et al. (2017). Risks and benefits of post-bariatric body contouring: A systematic review. *Journal of Plastic Surgery and Hand Surgery*, **51**(3): 168-175.
11. Teixeira, L. J., et al. (2019). Postoperative complications in body contouring surgery: A review of the literature. *American Journal of Surgery*, **218**(3): 477-483.
12. Goh, K. M., et al. (2019). Excess skin after bariatric surgery: A review of surgical options and outcomes. *Journal of Surgical Research*, **239**: 56-67.
13. Zhang, T., et al. (2018). Preoperative psychological evaluations and their impact on surgical outcomes in post-bariatric body contouring. *Aesthetic Surgery Journal*, **38**(6): 686-693.
14. Cheung, C., et al. (2020). The impact of excess skin on physical function after weight loss. *Surgery for Obesity and Related Diseases*, **16**(3): 441-448.
15. Zawawi, F. H., et al. (2021). The relationship between nutritional status and surgical outcomes in body contouring surgery. *Plastic and Reconstructive Surgery*, **148**(2): 332-339.
16. Santin, A., et al. (2019). The effect of preoperative weight and skin laxity on outcomes of body contouring surgery. *Aesthetic Plastic Surgery*, **43**(5): 1241-1248.
17. Feng, X., et al. (2018). The psychological impact of bariatric surgery on body image and quality of life. *Bariatric Surgical Practice and Patient Care*, **13**(3): 89-95.
18. Wu, Y., et al. (2020). Complications in post-bariatric body contouring surgery: A review of 1,000 cases. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, **73**(3): 526-533.
19. Khosravi, M. J., et al. (2020). Evaluation of patient-reported outcomes following body contouring surgery after weight loss. *Surgical Endoscopy*, **34**(6): 2435-2444.
20. Stowers, R. J., et al. (2020). Long-term outcomes of body contouring surgery after bariatric surgery: A retrospective study. *Obesity Surgery*, **30**(4): 1140-1149.

Dr. Adegu William, University of Nairobi, department of plastic, reconstructive and aesthetic surgery, Dr. Benjamin Wabwire, Kenyatta National Hospital, Dr. Protus Kituyi, Prof. Stanley Khainga, Moi University, Department of surgery and Anesthesiology and Prof. Ferdinand W. Nang'ole, University of Nairobi, Department of Plastic, Reconstructive and Aesthetic surgery

Corresponding Author: Dr. Adegu William, University of Nairobi, department of plastic, reconstructive and aesthetic surgery: Email address: jacobadegu@yahoo.com. P. O. Box 1663-50100, Kakamega, Kenya.

Disclaimer: This scientific communication was not peer reviewed and the opinions expressed herein are of the Authors only.