

Panfrican Journal of Plastic Reconstructive and Aesthetic Surgery Vol. 1 No. 2 December 2024

SENSATE CHIMERIC ANTEROLATERAL THIGH FLAP FOR HEAD AND NECK RECONSTRUCTION FOLLOWING CANCER EXTIRPATION IN A HIV POSITIVE PATIENT: CASE REPORT

Nyairo M.D, Kerubo L.J, Nyabuto C, Ogechi J, Kiriga M, Aswani J, Vilembwa A and Nangole F.W, Department of Surgery, University of Nairobi, P.O. Box 30197-00100, Nairobi, Kenya.

Corresponding Author: Dr. D. Nyairo, Department of Surgery, University of Nairobi, P.O. Box 30197-00100, Nairobi, Kenya.

SENSATE CHIMERIC ANTEROLATERAL THIGH FLAP FOR HEAD AND NECK RECONSTRUCTION FOLLOWING CANCER EXTIRPATION IN A HIV POSITIVE PATIENT: CASE REPORT

M. D. NYAIRO, L. J. KERUBO, C. NYABUTO, J. OGECHI, M. KIRIGA, J. ASWANI, A. VILEMBWA
and F. W. NANGOLE

SUMMARY

Radical tumor ablations in the head and neck area require intricate reconstructive procedures.

This is even more so in the setting of HIV where more advanced disease is observed at presentation resulting in poorer outcomes and increased complications. Free tissue transfer has enabled preservation of function and aesthetics while upholding oncologic surgery principles. The anterolateral thigh (ALT) flap is a very reliable workhorse flap in this regard.

We present a case of a 51-year-old HIV positive male with oral squamous cell carcinoma that was resected and the resultant defect reconstructed with a free, sensate, chimeric anterolateral thigh (ALT) flap. The patient was followed up closely. Initial intensive free flap monitoring and subsequent routine clinical reviews were undertaken. Six-week postoperative review revealed a healthy well incorporated flap that successfully resurfaced both intraoral and cutaneous defects.

A chimeric ALT flap is an excellent choice to resurface both the intraoral and cutaneous defects.

The lateral femoral cutaneous nerve of the thigh can be used to restore sensation in complex through and through cheek defects after oncologic surgery.

Key words: Chimeric, sensate flap, HIV positive, Head and neck reconstruction

INTRODUCTION

Head and neck cancer incidence is ranked third of all malignancies in developing nations (1) with oral squamous cell carcinoma being the highest at 40.6% (2,3).

HIV infection is associated with increased incidence, advanced disease and poorer outcomes of these malignancies which necessitates larger, more complex flap designs to achieve cover (4–11).

Chimeric ALT flaps are ideal for defects demanding reconstruction in greater than two dimensions. The lateral femoral cutaneous nerve of the thigh can be utilized for sensory restoration at the recipient site. We present a case of a HIV positive 51-year-old man in whom a sensate chimeric ALT flap was used to resurface a complex face defect after excision of advanced oral squamous cell carcinoma.

Case presentation

51-year-old HIV positive male on HAART for 14 years presented with an exophytic ulcerated oral squamous cell carcinoma involving right cheek and mandible.




Figure 1: Preoperative view of the mass.

The surgery involved right sub-mandibulectomy done at para symphysis level, disarticulation on the right and bilateral lymph node dissection.

The resultant cutaneous defect was approximately 16cm by 12cm and an intraoral defect of 8cm by 4cm.



Figure 2: Resultant through and through cheek defect

Fasciocutaneous ALT flap was raised with two musculocutaneous perforators then divided to match the defects, each segment with its own perforator.



Figure 3: (Left to right) the lateral femoral cutaneous nerve of the thigh, ALT flap being retracted, proximal perforator, distal perforator




Figure 4: Patient approximately 6 weeks post-operatively

DISCUSSION

Since its description by Song et al in 1984 (12), various authors have published on the versatility, vascular anatomic variations, surgical modifications and applications of the ALT flap in head and neck reconstruction (13–17).

ALT flap has good pliability and may be folded for the reconstruction of both the inner and outer lining of through and through cheek defects. The vascular pattern also allows the use of a more versatile design with double skin paddles based on multiple perforators (13,16). It may be raised with vastus lateralis muscle as a myocutaneous flap or combined with adjacent flaps according to the chimeric flap principle, to reconstruct large or complex 3-dimensional defects (17–19).

There is a dearth of knowledge on free flap surgery in patients living with HIV. Studies have reported higher surgical complication rates and flap loss in these patients implicating low CD4 counts, vasculitis, and sepsis as contributing factors (11,20,21). Few studies have demonstrated comparable complication rates between patients with and without HIV (20).

This case report shows that successful complex free flap reconstruction can be undertaken in the setting of HIV with proper patient selection.

CONCLUSION

The ALT flap is highly versatile and offers flexibility in resurfacing complex defects of the head and neck region. It can be raised with the lateral femoral cutaneous nerve of the thigh to restore protective sensation after oncologic tumor resection. Patients living with HIV can have successful microsurgical reconstruction with optimal patient selection and preparation.

REFERENCES

1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. 2015 Mar;**65**(2):87–108.
2. Onyango JF, Awange DO, Njiru A, Macharia IM. Pattern of occurrence of head and neck cancer presenting at Kenyatta National Hospital, Nairobi. *East Afr Med J*. 2006 May;**83**(5):288–91.
3. Gathere S, 1*, Mutuma G, 2, Korir A, 3, et al. Head and Neck Cancers four year trend at the Nairobi Cancer Registry. *Afr J Health Sci* 2011 1930–35.
4. Herida M, Mary-Krause M, Kaphan R, Cadranet J, Poizot-Martin I, Rabaud C, et al. Incidence of non-AIDS-defining cancers before and during the highly active antiretroviral therapy era in a cohort of human immunodeficiency virus-infected patients. *J Clin Oncol Off J Am Soc Clin Oncol*. 2003 Sep 15;**21**(18):3447–53.

5. McGinnis GJ, Ning MS, Bvochora-Nsingo M, Chiyapo S, Balang D, Ralefala T, et al. Management of Head and Neck Cancers With or Without Comorbid HIV Infection in Botswana. *The Laryngoscope*. 2021 May; **131**(5):E1558–66.
6. Salahuddin S, Cohen O, Wu M, Perez Irizarry J, Vega T, Gan G, et al. Human Immunodeficiency Virus Is Associated With Poor Overall Survival Among Patients With Head and Neck Cancer. *Clin Infect Dis Off Publ Infect Dis Soc Am*. 2023 Apr 17; **76**(8):1449–58.
7. Shiels MS, Pfeiffer RM, Gail MH, Hall HI, Li J, Chaturvedi AK, et al. Cancer burden in the HIV-infected population in the United States. *J Natl Cancer Inst*. 2011 May 4; **103**(9):753–62.
8. Deeken JF, Tjen-A-Looi A, Rudek MA, Okuliar C, Young M, Little RF, et al. The Rising Challenge of Non-AIDS-Defining Cancers in HIV-Infected Patients. *Clin Infect Dis Off Publ Infect Dis Soc Am*. 2012 Nov 1; **55**(9):1228–35.
9. Chiao EY, Coghill A, Kizub D, Fink V, Ndlovu N, Mazul A, et al. The effect of non-AIDS-defining cancers on people living with HIV. *Lancet Oncol*. 2021 Jun; **22**(6):e240–53.
10. Singh B, Balwally AN, Shaha AR, Rosenfeld RM, Har-El G, Lucente FE. Upper aerodigestive tract squamous cell carcinoma. The human immunodeficiency virus connection. *Arch Otolaryngol Head Neck Surg*. 1996 Jun; **122**(6):639–43.
11. Nangole WF, Khainga S, Aswani J, Kahoro L, Vilembwa A. Free Flaps in a Resource Constrained Environment: A Five-Year Experience – Outcomes and Lessons Learned. *Plast Surg Int*. 2015;2015:194174.
12. Song YG, Chen GZ, Song YL. The free thigh flap: a new free flap concept based on the septocutaneous artery. *Br J Plast Surg*.
13. Zaretski A, Wei FC, Lin CH, Cheng MH, Tsao CK, Wallace CG. Anterolateral Thigh Perforator Flaps in Head and Neck Reconstruction. *Semin Plast Surg*. 2006 May; **20**(2):64–72.
14. Shieh SJ, Chiu HY, Yu JC, Pan SC, Tsai ST, Shen CL. Free anterolateral thigh flap for reconstruction of head and neck defects following cancer ablation. *Plast Reconstr Surg*. 2000 Jun; **105**(7):2349–57; discussion 2358–2360.
15. Koshima I. Free Anterolateral Thigh Flap for Reconstruction of Head and Neck Defects following Cancer Ablation. *Plast Reconstr Surg*. 2000 Jun; **105**(7):2358–60.
16. Chana JS, Wei F chan. A review of the advantages of the anterolateral thigh flap in head and neck reconstruction. *Br J Plast Surg*. 2004 Oct 1; **57**(7):603–9.
17. Huang WC, Chen HC, Jain V, Kilda M, Lin YD, Cheng MH, et al. Reconstruction of Through-and-Through Cheek Defects Involving the Oral Commissure, Using Chimeric Flaps from the Thigh Lateral Femoral Circumflex System. *Plast Reconstr Surg*. 2002 Feb; **109**(2):433.
18. Koshima I, Yamamoto H, Hosoda M, Moriguchi T, Orita Y, Nagayama H. Free combined composite flaps using the lateral circumflex femoral system for repair of massive defects of the head and neck regions: an introduction to the chimeric flap principle. *Plast Reconstr Surg*. 1993 Sep; **92**(3):411–20.
19. Wei F chan, Jain V, Celik N, Chen H chi, Chuang DCC, Lin Chung. Have We Found an Ideal Soft-Tissue Flap? An Experience with 672 Anterolateral Thigh Flaps. *Plast Reconstr Surg*. 2002 Jun; **109**(7):2219.
20. Runodada PM, Chihaka OB, Muguti GI. Surgical outcomes in HIV positive patients following major surgery at two tertiary institutions in Harare, Zimbabwe. *Int J Surg Open*. 2020 Jan 1; **22**:12–7.
21. Brand M, Woodiwiss AJ, Michel F, Nayler S, Veller MG, Norton GR. Large vessel adventitial vasculitis characterizes patients with critical lower limb ischemia with as compared to without human immunodeficiency virus infection. *PloS One*. 2014; **9**(8):e106205.